

Scoping Review

Isokinetic Dynamometry for External and Internal Rotation Shoulder Strength in Youth Athletes: A Scoping Review

Ian Leahy^{1a}, Erin Florkiewicz¹, Mary P. Shotwell²

¹ Orthopedics, Rocky Mountain University of Health Professions, ² Integrative Health Sciences, Rocky Mountain University of Health Professions

Keywords: Isokinetic dynamometry, shoulder, youth, overhead athlete, external rotation, internal rotation, strength, peak torque

<https://doi.org/10.26603/001c.125765>

International Journal of Sports Physical Therapy

Vol. 19, Issue 12, 2024

Background

Accurately measuring shoulder strength in overhead athletes is critical, as sufficient strength is essential for safe and sustained performance during repetitive athletic movements. Isokinetic dynamometry (ID) offers dynamic strength assessments that surpass the capabilities of static methods, such as manual muscle testing and handheld dynamometry. The dynamic assessment provided by ID may enhance upper extremity evaluation, aiding in the prediction of injury risk and the determination of return-to-sport criteria for overhead athletes.

Purpose

The purpose of this review was to examine the existing literature concerning the application of isokinetic shoulder strength testing in rehabilitation and clinical decision-making processes among youth athletes who perform repetitive overhead activities.

Study Design

Scoping review

Methods

A comprehensive literature search was conducted using PubMed and EBSCO Host databases, covering publications from 2000-2024. Search terms included “isokinetic dynamometry,” “shoulder,” and “youth athlete.” Inclusion criteria focused on youth athletes (<18 years) engaged in overhead sports, excluding those with neurological conditions or those designated as college or professional athletes. The PRISMA-ScR guidelines were followed.

Results

A total of 23 articles met the inclusion criteria. Volleyball and swimming were the most studied sports, with the most common testing position being the seated 90/90 position. Variations in testing speeds and outcome measures, such as peak torque and external rotation (ER) ratios, were identified.

Conclusions

Isokinetic dynamometry is a valuable tool for assessing shoulder strength in youth overhead athletes. It provides critical insights into muscle strength dynamics, aiding in injury prevention and rehabilitation. Further research is needed to optimize strength

^a Corresponding Author:

Ian Leahy, PT, DPT, OCS, SCS, CSCS, FAAOMPT
577 Washington Crossing Rd, Newtown PA 18940
267-226-5360
Leahydpt@gmail.com

assessment protocols and enhance clinical decision-making for safe return-to-sport practices.

BACKGROUND

Within a larger framework in the decision process for return to sports, recommendations for overhead athletes are reported considering normalized shoulder strength and scapular mechanics, range of motion (ROM), and successful completion of a plyometric program.¹ However, many clinicians find it challenging to accurately measure shoulder strength due to limitations in equipment and expertise.

Shoulder strength assessment generally consists of isometric testing in neutral positions via manual muscle testing (MMT), handheld dynamometry (HHD), or isokinetic dynamometry (ID). Normative values for external rotation (ER) and internal rotation (IR) strength using HHD has been obtained in healthy adults,^{2,3} and physically active collegiate male and females.⁴ Studies indicate that HHD provides an assessment of shoulder IR/ER strength assessment that is highly correlated to assessments performed with ID.⁵ The concern, however, with MMT and HHD testing is that reliability of these methods is highly dependent on clinician size and strength.⁶ Additionally, because the testing position is static, HHD lacks the assessment of dynamic components associated with upper extremity movements required during overhead sports.⁷ In addition to torque output as a measure of strength, ID can capture endurance deficits and strength ratio imbalances which are predictors of shoulder injury.⁸ Ellenbecker demonstrated a significant difference in bilateral IR and ER strength measures between MMT grades and isokinetic dynamometry, widely recognized as the gold standard for measuring muscle strength and muscle endurance.⁹

Understanding the balance between the strength of agonist and antagonist muscles is crucial in evaluating and rehabilitating overhead athletes due to the intricate muscular activation patterns necessary for stability of the glenohumeral joint.⁷ One strategy for strength testing of the shoulder in multiple positions throughout the range of motion is ID. Multiple authors have reported normative isokinetic strength assessment data for athletes involved in judo,¹⁰ tennis,^{11,12} badminton,^{13,14} and volleyball.¹⁵ In baseball, isokinetic strength testing has been performed in high school athletes,¹⁶ collegiate athletes,^{17,18} and professional athletes,¹⁹ yet limited information is available regarding normative values of isokinetic shoulder strength in youth athletes.

Gaps remain in the literature, particularly regarding the establishment of normative values for younger, skeletally immature athletes and how these values differ across various stages of development. The purpose of this review was to examine the existing literature concerning the application of isokinetic shoulder strength testing in rehabilitation and clinical decision-making processes among youth athletes who perform repetitive overhead activities.




Figure 1. PRISMA flow diagram showing the literature search, screening, and eligibility results.

METHODS

This literature review was conducted in accordance with the recommendations of the "Preferred Reporting Items for Systematic Reviews and Meta Analysis extension for Scoping Reviews (PRISMA-ScR).²⁰

SEARCH STRATEGY

The literature review was performed with the databases PubMed, and EBSCO Host, which provides a range of databases, e-journals and e-books. Search terms included in the search strategies included phrases such as: "isokinetic dynamometry AND shoulder," "isokinetic dynamometry AND shoulder AND youth athlete." (See Appendix A for full search criteria). The search was limited to full-text available human research published between 2000-2024, including participants ages <18 years associated with an organized, overhead sport, and published in the English language. Exclusion criteria were articles pertaining to any neurological condition, and athletes designated as collegiate or professional.

RESULTS

Electronic database searches identified 475 total studies. The scoping review included 23 studies²¹⁻⁴³ after duplicate removal, title/abstract screen, and full text review.

The PRISMA diagram (Figure 1) outlines the search process in its entirety.

Table 1. Use of Isokinetic Dynamometry (ID) in Various Sports and ID Equipment Parameters

Author	Sample / (Mean Age)	Sport	Type of Dynamometer	Position
Batalha N ²¹	25 males (13.28 yrs)	Swimming	Biodex System-3	Seated 90/90 position
Batalha NM ²⁴	36 males (14.25 yrs)	Swimming	Biodex System-3	Seated 90/90 position
Batalha N ²³	40 males (14.65 yrs)	Swimming	Biodex System-3	Seated 90/90 position
Batalha NM ²⁵	40 males (14.65 yrs)	Swimming	Biodex System-3	Seated 90/90
Batalha N ²²	49 males (14.48 yrs)	Swimming	Biodex System-3	Seated 90/90
Clements AS ²⁶	18 males (13-16 yrs)	Baseball	Biodex Multi-Joint Dynamometer	Seated 90/90
de Lira CAB ²⁷	28 males (15.5 yrs)	Volleyball	Biodex System-3	Seated (angle not specified)
Dupuis, C ²⁸	10 males (15.87 yrs)	Baseball	Kin-Com	Seated 90/90
Duzgun I ²⁹	24 athletes (14.5 yrs)	Volleyball	Isomed 2000 Dynamometer	Seated 90/90 position
Eshghi S ³⁰	32 males (17.5 yrs)	Volleyball	Biodex System-4	Seated position 90 degrees abduction / 30 degrees shoulder flexion
Guney H ³¹	65 athletes (16.1 yrs)	Volleyball & Basketball	Isomed2000 D&R	Seated 90/90
Lee DR ³²	23 males (18.2 yrs)	Baseball	Biodex Isokinetic Machine	Seated 90/90 position
Mascarin NC ³³	26 females (15.3 yrs)	Handball	Biodex Isokinetic Machine	Not Discussed
Mascarin NC ³⁴	39 females (15.3 yrs)	Handball	Biodex Isokinetic Dynamometer	Seated 90/90
Mickevičius M ³⁵	14 boys (11-12 yrs)	Baseball	Biodex System-3	Seated 90/90
Mohamed IW ³⁶	16 males (14.8 yrs)	Weightlifting	Multi Joint System-3 Pro	Seated 45/90 position Repeat testing done @ 90/90
Mulligan IJ ³⁷	39 males (15.36 yrs)	Baseball	Kin-Com Dynamometer	Seated: 90 degrees elbow flexion / 30 degrees abduction in scapular plane
Pawlik D ³⁸	12 females (12-13 yrs)	Volleyball	Multi-joint 4 Dynamometer Biodex	Seated 90/90
Pontaga J ³⁹	14 males (14.6 yrs)	Handball	REV-9000	Seated 90/90 position
Saccol MF ⁴⁰	40 athletes (14 yrs)	Tennis	Cybex-6000	Supine 90/90
van Cingel R ⁴¹	40 females (17.6 yrs)	Handball	2-Humac Norm dynamometer	Supine 90/90
Vodička T ⁴²	20 males (13.23 yrs)	Tennis	Human Norm CSMI	Supine 90/90 position
Yildiz Y ⁴³	40 males (17.2 yrs)	Volleyball, Handball, Tennis	Cybex Norm	Supine 90/90

ISOKINETIC DYNAMOMETRY (ID) IN YOUTH REPETITIVE OVERHEAD ATHLETES

Of the included studies, volleyball was the most common overhead sport where ID was used for assessment.^{27,29-31, 38,43} The second most common sport was swimming²¹⁻²⁵ followed by handball,^{33,34,39,41,43} baseball,^{26,28,32,37} tennis,^{40,42,43} weightlifting,³⁶ and basketball.³¹ In the articles that fit the inclusion criteria, two pertained to symptomatic patients,^{31,35} whereas the remaining articles looked at isokinetic normative values, or the effects of repetitive sport specific movements. Table 1 describes the use of ID in various sports along with the variety of parameters used for testing ID.

ISOKINETIC TESTING POSITION

Testing positions throughout studies varied, but trends are seen when testing the upper extremity (UE) of youth athletes with ID as illustrated in Table 1. The most utilized testing position for shoulder IR/ER is the seated position with the shoulder abducted to 90 degrees and the elbow flexed to 90 degrees, commonly known as the 90/90 position. It is postulated that this position, more than the arm at a neutral position, specifically addresses muscle function often required for an overhead athlete.⁴⁴ The use of the 90/90 position also demonstrates strong intraclass coefficient for test/re-test reliability.⁴⁵ Ellenbecker has discussed using a modified position of 30 degrees shoulder abduction, 30 degrees shoulder forward flexion, and 30 degrees diago-

nal tilt of the dynamometer to determine tolerance of the UE strength test prior to using the ID at a 90/90 position.⁷ Additional alterations to the 90/90 positions used in research are the seated 90/30 (90 degrees elbow flexion and 30 degrees shoulder abduction),³⁷ seated 90/45 (90 degrees elbow flexion and 45 degrees shoulder abduction),³⁶ seated 90/30 with 90 degrees of shoulder abduction and 30 degrees of shoulder flexion,³⁰ and the 90/90 supine position.⁴⁰⁻⁴³

SPEEDS USED FOR TESTING

[Table 2](#) provides information regarding the variations in the speeds used for testing youth athletes. Most studies assessed participants at multiple speeds for different outcome measures or for different sports-related contexts. Common speeds for assessment of isokinetic peak torque both concentrically and eccentrically include 60 deg/sec,^{22-25,27,29,30,32-34,38-41} 90 deg/sec,^{28,31,37,39,43} 120 deg/sec,^{26,36,41} 180 deg/sec,^{21-25,28-30,32,37,38,40,42} 240 deg/sec,^{27,33,34,36,39} 300 deg/sec,^{38,42} and 360 deg/sec.³⁶

OBJECTIVE MEASURES FROM ISOKINETIC DYNAMOMETRY

PEAK TORQUE

In many studies pertaining to the use of ID in youth athletes, peak torque, normalized via body weight, was the primary objective measure.^{26,27,29,40} Peak torque is then used to compute measures of bilateral symmetry, using left and right values, and unilateral strength ratios using shoulder external and internal rotation strength values. Research indicates that repetitive sports-related movements lead to discernible discrepancies in peak torque between shoulder external rotation (ER) and internal rotation (IR) strength when comparing the dominant and non-dominant arms in athletes. Repetitive motions during youth sports such as volleyball,^{27,38,43} swimming,²¹⁻²⁵ and tennis⁴⁰ result in increased IR peak torque in the dominant arm of the athlete. Variations exist among age groups within the same sport as evidenced by youth handball athletes demonstrating higher ER and IR peak torque in the dominant arm compared to non-dominant arm in athletes with a mean age of 17.6 years.⁴¹ Conversely, younger players (mean age of 14.6 years) showed no statistically significant difference between ER and IR peak torque.³⁹ In a study of healthy teenage baseball players, Dupuis found greater concentric ER strength in the dominant arm at speeds of 90°/sec and 180°/sec.²⁸ In contrast, Mickevičius observed decreased ER strength among younger baseball players (average age 11.6 years) who had a history of shoulder pain.³⁵ It has been hypothesized that differences in peak torque output results in higher likelihood of overuse type injuries in youth athletes.^{24,27,41}

ER:IR RATIO

Eight studies^{21-25,35,40,42} used concentric external rotation to concentric internal rotation strength ratios during for their objective measurements whereas six studies^{28,30,31,34,}

^{41,43} investigated various forms of eccentric to concentric ratios between ER and IR.

Comparison of shoulder agonist and antagonist muscle groups helps identify specific muscular imbalances.⁷ Variations in the dominant arm's external rotation to internal rotation (ER:IR) ratios are frequently observed, particularly in cases where imbalances occur in the shoulder's agonist-antagonist muscle relationships. These imbalances can signify weaknesses that heighten athletes' vulnerability to injury.^{24,40} Ideal ER:IR has been defined as 66%, which remains constant throughout the velocity spectrum⁴⁶; however, this ratio is described in skeletally mature athletes.⁷

A common theme in youth overhead athletes is decreased ER:IR ratio in the dominant arm^{21-24,34,40-42} due to the theorized increase in IR peak torque because of sport specific repetitive motion. Although these imbalances do not result in changes in athletic performance, there is fear that if not addressed in the younger athlete they can result in overuse type injury.⁴⁰

There are unique variations of the ER:IR used in ID assessment of youth athletes as evidenced by discussion of ER:IR acceleration and ER:IR deceleration described by Yildiz in youth volleyball players.⁴³ The concept eccentric to concentric ER:IR is defined as the ratio of balance for muscle activity where the shoulder medial rotators eccentrically control external rotation, while lateral rotators eccentrically control internal rotation, ensuring optimal shoulder function.⁴⁷ This explanation is similar to Dupuis' definition of Dynamic Control Ratio (DCR) in youth baseball players.²⁸ Both Yildiz and Dupuis investigate the eccentric and concentric relationship of the ER:IR ratio of the shoulder. In each case, the ER:IR ratio into the acceleration phase (concentric ER to eccentric IR) was observed to be higher in the dominant arm,^{28,43} whereas the ER:IR ratio into the deceleration phase (eccentric ER to concentric IR) is lower in the dominant arm of the youth volleyball athlete,⁴³ but no difference existed in the youth baseball athlete.²⁸

DISCUSSION

This scoping review highlights the use of isokinetic dynamometry of the youth overhead athlete to obtain multiple objective measures. In adolescent overhead athletes, the repetitive nature of sport-specific movements imposes unique demands on the dynamic muscular control of the glenohumeral joint. This demand is highlighted by fluctuations in the external rotation to internal rotation peak torque ratio, a phenomenon hypothesized to significantly impact the vulnerability of the shoulder complex to injury.²¹⁻²⁵ Various studies support this hypothesis, including symptomatic athletes³⁵ and those with glenohumeral internal rotation deficits.³¹

In sports like baseball, dynamic muscular control of the glenohumeral joint is essential, particularly given the substantial stress experienced across the upper extremity joints during pitching. The angular velocities of shoulder internal rotation and elbow extension can range from 1,000

Table 2. Isokinetic Dynamometry Testing Speed and Other Objective Measures

Author	Testing Speed	Objective Measures
Batalha N ²¹	60 deg/sec 180 deg/sec	- PT IR - PT ER - ER:IR ratio
Batalha N ²⁴	60 deg/sec 180 deg/sec	- PT - ER/IR ratio
Batalha N ²³	60 deg/sec 180 deg/sec	- Max torque - ER/IR ratio - norm is 66-75%
Batalha N ²⁵	60 deg/sec 180 deg/sec	- PT - ER/IR ratios
Batalha N ²²	60 deg/sec 180 deg/sec	- PT - ER/IR ratio
Clements AS ²⁶	120 deg/sec	PT/BW
de Lira CAB ²⁷	60 deg/sec 240 deg/sec	- Absolute PT - Relative PT (divided by BW) - Absolute total work - Relative total work (divided by BW) - Conventional strength ratio - Functional strength ratio
Dupuis, C ²⁸	90 deg/sec 180 deg/sec	- PT - IR con / ER ecc (performance ratio) - ER con / IR ecc (cocking ratio)
Duzgun I ²⁹	60 deg/sec 180 deg/sec	- PT / BW - Total work
Eshghi S ³⁰	60 deg/sec 180 deg/sec	- FDR - EReccn:IRcon
Guney H ³¹	90 deg/sec	- ER:IR - Max eccentric ER torque / Max concentric IR torque
Lee D ³²	60 deg/sec 180 deg/sec	PT
Mascarin N ³³	60 deg/sec 240 deg/sec	PT and total work
Mascarin N ³⁴	60 deg/sec 240 deg/sec	- at 60 deg/sec: - IR PT - CR (ER con / IR con) - at 240 deg/sec - IR PT - IR avg power - FR (ER ecc / IR con) - ER PT was measured at both speeds
Mickevičius M ³⁵	120 deg/sec	- PT - ER:IR ratio
Mohamed IW ³⁶	120 deg/sec 240 deg/sec 360 deg/sec	- Relative PT - Time to PT
Mulligan I ³⁷	90 deg/sec 180 deg/sec	PT and Total work for both movements at both speeds
Pawlik D ³⁸	60 deg/sec 180 deg/sec 300 deg/sec	Concentric IR PT
Pontaga I ³⁹	60 deg/sec 90 deg/sec	- PT - Avg power

Author	Testing Speed	Objective Measures
	240 deg/sec	
Saccol M ⁴⁰	60 deg/sec 180 deg/sec	- PT / BW - Total work / BW - ER/IR ratio
Vodička T ⁴²	180 deg/sec 300 deg/sec	- Peak torque - ER/IR ratio
Yildiz Y ⁴³	90 deg/sec	- ER ecc / IR con - ER con / IR ecc - Both in the late cocking phase of overhead motion and the deceleration phase

Key: PT= peak torque, ER= external rotation, IR= internal rotation, BW= body weight, CR= cocking ratio, FDR= functional deceleration ratio, FR= functional ratio, DCR= dynamic control ratio, CON = concentric, ECC = eccentric

deg/sec to 7,200 deg/sec in youth and collegiate pitchers, respectively, from maximum external rotation through the acceleration phase.^{48,49} While ID testing velocities of 240 deg/sec do not fully replicate these speeds, they still yield valuable data on shoulder musculature function at higher isokinetic speeds.⁵⁰ [Table 3](#) provides a detailed breakdown of these different sport-specific findings.

GAPS IN THE LITERATURE

ID studies of youth athletes skew towards specific sports such as volleyball and swimming. Limited information is available regarding the use of ID for assessment of athletes participating in youth baseball. In reviewing the ages of participants recruited for ID articles in youth athletes, the youngest subjects were 11 years old. However, pitching and repetitive throwing in baseball often begins as early as 6-7 years of age. Studies included in a systematic review suggest that starting repetitive pitching at 10 years of age or younger increases the risk of upper extremity-related injury.⁵¹ In most sports, children are playing and specializing in sport positions at a much younger age than captured in this review, but changes in shoulder IR strength between dominant and non-dominant arm has been seen in youth throwers under the age of 10 using HHD.⁵² Understanding dynamic strength values and muscular balance relationships with measurements from ID in skeletally immature athletes and the changes that occur with repetitive overhead motions can assist in developing injury prevention, rehabilitation programs, and return to sport decision making for youth throwing athletes.

LIMITATIONS

Only studies in English languages were included in the review and the range of articles searched were between 2000-2024. Also, no quality assessment of the studies was performed, which may limit the impact of the findings.

CONCLUSION

Isokinetic dynamometry as an assessment in youth overhead athletes provides insights into shoulder strength

(torque) and strength ratios. There is variety in the assessment parameters utilized in the included studies of youth athletes in various sports. By precisely evaluating isokinetic strength, clinicians can gauge the shoulder's dynamic control, facilitating the tailored design of training programs. Isokinetic assessments allow for detailed analysis of power and the balance between agonist and antagonist muscle groups during movement, enabling the identification of specific strength impairments and guiding targeted interventions to enhance upper extremity loading tolerance for repetitive overhead activities.

CONFLICT OF INTEREST

The authors declare that they have no conflicts of interest related to the content of this manuscript.

Submitted: June 19, 2024 CDT. Accepted: September 17, 2024 CDT. Published: December 01, 2024 CDT.

© The Author(s)

Table 3. Sports Specific Isokinetic Dynamometry (ID) Strength Findings

SPORT	STUDY	ID SPORT SPECIFIC STRENGTH OUTCOMES
Volleyball	De Lira ²⁷	Higher IR peak torque in dominant arm
	Duzgun ²⁹	ER strength increased with weighted jump rope in dominant arm
	Esghi ³⁰	- Functional Deceleration Ratio (eccentric to concentric strength of IR and ER) in dominant arm - Increased ER:IR ratio after strengthening program
	Pawlik ³⁸	Higher IR peak torque in dominant arm
Swimming	Batalha ²¹	Higher IR peak torque in dominant arm
	Batalha ²²	- Decreased ER:IR ratio in dominant arm in swimmers vs control - Higher IR peak torque in swimmer dominant arm
	Batalha ²³	Decreased ER:IR ratio in dominant arm for swim only group
	Batalha ²⁴	- Decreased ER:IR in dominant arm for swimmers- - Higher IR peak torque in dominant arm for swimmers
	Batalha ²⁵	- Decreased ER:IR in dominant arm for swimmers - Training improved ER peak torque in dominant arm
Baseball	Clements ²⁶	Peak torque of dominant arm not associated with throwing velocity
	Dupuis ²⁸	- Higher ER peak torque in dominant arm - ER:IR lower in dominant arm - ER concentric :IR eccentric in cocking phase was higher in dominant arm - ER eccentric:IR concentric during deceleration was no different side to side
	Lee ³²	IR and ER peak torque increased with CKC strengthening in dominant arm
	Mickevicius ³⁵	- ER eccentric strength was weaker in baseball players dominant arm - Decreased ER:IR in dominant arm for baseball players with pain - IR eccentric: IR concentric – lower in dominant arm - ER eccentric: IR concentric – lower in dominant arm - ER concentric: IR eccentric – lower in dominant arm
	Mulligan ³⁷	- Decreased ER:IR ratio in the dominant arm - No significant difference between eccentric and concentric peak torque between dominant vs non-dominant arm
Handball	Mascarin ³³	- ER:IR lower in dominant arm - ER eccentric : IR concentric was lower in dominant arm - ER:IR ratio unchanged in dominant arm with strength training
	Mascarin ³⁴	- All participants demonstrated weak IR of the dominant arm - ER:IR – 82.8 in experimental group vs 77.6 in control group - Strength testing did not result in improvement in IR strength - Strength training did not improve strength ratios - Strength training improved average power
	Pontaga ³⁹	- No statistically significant differences between dominant and non-dominant arm peak torque - Average power of IR was higher in dominant arm
	Van Cingel ⁴¹	- ER:IR lower in dominant arm of handball players - Higher ER and IR peak torque in dominant arm
Tennis	Sacco ⁴⁰	- Decreased ER:IR on dominant arm - Higher ER strength in youth male tennis players - Higher IR strength on dominant arm for both youth male and female athletes
	Vodička ⁴²	- Decreased ER:IR ratio on dominant arm - Higher ER strength in dominant arm - Higher IR strength on dominant side
Weightlifting	Mohamed ³⁶	- ER:IR ratio of dominant arm improved with isokinetic strength program - ER:IR ratio of dominant arm improved with isotonic strength program - IR and ER peak torque of dominant arm increased in both training groups

Key: ER= external rotation, IR= internal rotation



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

REFERENCES

1. Sgroi TA, Zajac JM. Return to throwing after shoulder or elbow injury. *Curr Rev Musculoskelet Med*. 2018;11(1):12-18. doi:[10.1007/s12178-018-9454-7](https://doi.org/10.1007/s12178-018-9454-7)
2. Bradley H, Pierpoint L. Normative values of isometric shoulder strength among healthy adults. *Int J Sports Phys Ther*. 2023;18(4):977-988. doi:[10.26603/001c.83938](https://doi.org/10.26603/001c.83938)
3. Harbin G, Leyh C, Harbin A. Upper extremity strength: Normative reference data among uninjured employees. *Work*. 2020;67(4):979-991. doi:[10.3233/wor-203348](https://doi.org/10.3233/wor-203348)
4. Westrick RB, Duffey ML, Cameron KL, Gerber JP, Owens BD. Isometric shoulder strength reference values for physically active collegiate males and females. *Sports Health*. 2013;5(1):17-21. doi:[10.1177/1941738112456280](https://doi.org/10.1177/1941738112456280)
5. Chamorro C, Arancibia M, Trigo B, Arias-Poblete L, Jerez-Mayorga D. Absolute reliability and concurrent validity of hand-held dynamometry in shoulder rotator strength assessment: systematic review and meta-analysis. *Int J Environ Res Public Health*. 2021;18(17). doi:[10.3390/ijerph18179293](https://doi.org/10.3390/ijerph18179293)
6. Nicholas JA, Sapega A, Kraus H, Webb JN. Factors influencing manual muscle tests in physical therapy. *J Bone Joint Surg Am*. 1978;60(2):186-190. doi:[10.2106/00004623-197860020-00008](https://doi.org/10.2106/00004623-197860020-00008)
7. Ellenbecker TS, Davies GJ. The application of isokinetics in testing and rehabilitation of the shoulder complex. *J Athl Train*. 2000;35(3):338-350.
8. Bagordo A, Ciletti K, Kemp-Smith K, Simas V, Climstein M, Furness J. Isokinetic dynamometry as a tool to predict shoulder injury in an overhead athlete population: A systematic review. *Sports*. 2020;8(9). doi:[10.3390/sports8090124](https://doi.org/10.3390/sports8090124)
9. Ellenbecker TS. Muscular strength relationship between normal grade manual muscle testing and isokinetic measurement of the shoulder internal and external rotators. *Isokin Exerc Sci*. 1996;6:51-56. doi:[10.3233/IES-1996-6109](https://doi.org/10.3233/IES-1996-6109)
10. Marcondes FB, Castropil W, Schor B, Miana A, Vasconcelos R, Etchebehere M. Shoulder isokinetic performance in healthy professional judo athletes: normative data. *Acta Ortop Bras*. 2019;27(6):308-312. doi:[10.1590/1413-785220192706223708](https://doi.org/10.1590/1413-785220192706223708)
11. Niederbracht Y, Shim AL. Concentric internal and eccentric external fatigue resistanc of the shoulder rotator muscles in female tennis players. *N Am J Sports Phys Ther*. 2008;3(2):89-94.
12. Ellenbecker T, Roetert EP. Age specific isokinetic glenohumeral internal and external rotation strength in elite junior tennis players. *J Sci Med Sport*. 2003;6(1):63-70. doi:[10.1016/s1440-2440\(03\)80009-9](https://doi.org/10.1016/s1440-2440(03)80009-9)
13. Zulfikri N, Selvanayagam VS, Yusof A. Evaluation of shoulder and knee isokinetic strength profile among elite adolescent badminton players. *J Sport Rehabil*. 2021;30(5):717-724. doi:[10.1123/jsr.2019-0483](https://doi.org/10.1123/jsr.2019-0483)
14. Ng GY, Lam PC. A study of antagonist/agonist isokinetic work ratios of shoulder rotators in men who play badminton. *J Orthop Sports Phys Ther*. 2002;32(8):399-404. doi:[10.2519/jospt.2002.32.8.399](https://doi.org/10.2519/jospt.2002.32.8.399)
15. Hadzic V, Sattler T, Veselko M, Markovic G, Dervisevic E. Strength asymmetry of the shoulders in elite volleyball players. *J Athl Train*. 2014;49(3):338-344. doi:[10.4085/1062-6050-49.2.05](https://doi.org/10.4085/1062-6050-49.2.05)
16. Trunt A, Fisher BT, MacFadden LN. Athletic shoulder test differences exist bilaterally in healthy pitchers. *Int J Sports Phys Ther*. 2022;17(4):715-723. doi:[10.26603/001c.35722](https://doi.org/10.26603/001c.35722)
17. Lin HT, Ko HT, Lee KC, Chen YC, Wang DC. The changes in shoulder rotation strength ratio for various shoulder positions and speeds in the scapular plane between baseball players and non-players. *J Phys Ther Sci*. 2015;27(5):1559-1563. doi:[10.1589/jpts.27.1559](https://doi.org/10.1589/jpts.27.1559)
18. Dodds FT, Knotts SS, Penrod MI, Scoggins WA, Conners RT. Shoulder strength and range of motion between collegiate pitchers and position players in baseball. *Int J Exerc Sci*. 2020;13(6):123-130.
19. Kim BG, Lim SK, Kong S. The Relationship between scapular upward rotation and shoulder internal and external rotation isokinetic strength in professional baseball pitchers. *Healthcare*. 2021;9(6). doi:[10.3390/healthcare9060759](https://doi.org/10.3390/healthcare9060759)
20. Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. *Ann Intern Med*. 2018;169(7):467-473. doi:[10.7326/m18-0850](https://doi.org/10.7326/m18-0850)

21. Batalha N, Dias S, Marinho DA, Parraca JA. The effectiveness of land and water based resistance training on shoulder rotator cuff strength and balance of youth swimmers. *J Hum Kinet.* 2018;62:91-102. doi:[10.1515/hukin-2017-0161](https://doi.org/10.1515/hukin-2017-0161)
22. Batalha N, Marmeleira J, Garrido N, Silva AJ. Does a water-training macrocycle really create imbalances in swimmers' shoulder rotator muscles? *Eur J Sport Sci.* 2015;15(2):167-172. doi:[10.1080/17461391.2014.908957](https://doi.org/10.1080/17461391.2014.908957)
23. Batalha N, Raimundo A, Tomas-Carus P, Paulo J, Simão R, Silva AJ. Does a land-based compensatory strength-training programme influences the rotator cuff balance of young competitive swimmers? *Eur J Sport Sci.* 2015;15(8):764-772. doi:[10.1080/17461391.2015.1051132](https://doi.org/10.1080/17461391.2015.1051132)
24. Batalha NM, Raimundo AM, Tomas-Carus P, Barbosa TM, Silva AJ. Shoulder rotator cuff balance, strength, and endurance in young swimmers during a competitive season. *J Strength Cond Res.* 2013;27(9):2562-2568. doi:[10.1519/JSC.0b013e31827fd849](https://doi.org/10.1519/JSC.0b013e31827fd849)
25. Batalha NM, Raimundo AM, Tomas-Carus P, Marques MA, Silva AJ. Does an in-season detraining period affect the shoulder rotator cuff strength and balance of young swimmers? *J Strength Cond Res.* 2014;28(7):2054-2062. doi:[10.1519/jsc.0000000000000351](https://doi.org/10.1519/jsc.0000000000000351)
26. Clements AS, Ginn KA, Henley E. Correlation between muscle strength and throwing speed in adolescent baseball players. *Phys Ther Sport.* 2001;2(3):123-131. doi:[10.1054/ptsp.2000.0025](https://doi.org/10.1054/ptsp.2000.0025)
27. de Lira CAB, Vargas VZ, Vancini RL, Andrade MS. Profiling isokinetic strength of shoulder rotator muscles in adolescent asymptomatic male volleyball players. *Sports.* 2019;7(2). doi:[10.3390/sports7020049](https://doi.org/10.3390/sports7020049)
28. Dupuis C, Tourny-Chollet C, Delarue Y, Beuret-Blanquart F. Influence of baseball practice on strength ratios in shoulder rotator muscles: A new position for isokinetic assessment. *Isokin Exerc Sci.* 2004;12(2):149-157. doi:[10.3233/IES-2004-0167](https://doi.org/10.3233/IES-2004-0167)
29. Duzgun I, Baltaci G, Colakoglu F, Tunay VB, Ozer D. The effects of jump-rope training on shoulder isokinetic strength in adolescent volleyball players. *J Sport Rehabil.* 2010;19(2):184-199. doi:[10.1123/jsr.19.2.184](https://doi.org/10.1123/jsr.19.2.184)
30. Eshghi S, Zarei M, Abbasi H, Alizadeh S. The effect of shoulder injury prevention program on shoulder isokinetic strength in young male volleyball players. *Res Sports Med.* 2022;30(2):203-214. doi:[10.1080/15438627.2020.1860050](https://doi.org/10.1080/15438627.2020.1860050)
31. Guney H, Harput G, Colakoglu F, Baltaci G. The effect of glenohumeral internal-rotation deficit on functional rotator-strength ratio in adolescent overhead athletes. *J Sport Rehabil.* 2016;25(1):52-57. doi:[10.1123/jsr.2014-0260](https://doi.org/10.1123/jsr.2014-0260)
32. Lee DR, Kim LJ. Internal- and external-rotation peak torque in little league baseball players with subacromial impingement syndrome: improved by closed kinetic chain shoulder training. *J Sport Rehabil.* 2016;25(3):263-265. doi:[10.1123/jsr.2014-0333](https://doi.org/10.1123/jsr.2014-0333)
33. Mascarín NC, de Lira CAB, Vancini RL, da Silva AC, Andrade MS. The effects of preventive rubber band training on shoulder joint imbalance and throwing performance in handball players: A randomized and prospective study. *J Bodyw Mov Ther.* 2017;21(4):1017-1023. doi:[10.1016/j.jbmt.2017.01.003](https://doi.org/10.1016/j.jbmt.2017.01.003)
34. Mascarín NC, de Lira CAB, Vancini RL, de Castro Pochini A, da Silva AC, Dos Santos Andrade M. Strength training using elastic bands: improvement of muscle power and throwing performance in young female handball players. *J Sport Rehabil.* 2017;26(3):245-252. doi:[10.1123/jsr.2015-0153](https://doi.org/10.1123/jsr.2015-0153)
35. Mickevičius M, Rutkauskas S, Sipavičienė S, et al. Absence of bilateral differences in child baseball players with throwing-related pain. *Int J Sports Med.* 2016;37(12):952-957. doi:[10.1055/s-0042-106297](https://doi.org/10.1055/s-0042-106297)
36. Mohamed IW, Rahim MFA, Shaharudin S. Effects of isokinetic versus isotonic training on strength, power and muscular balance of rotator cuff muscles among advanced level of adolescent weightlifters. *Int J Appl Sports Sci.* 2017;29(2):143-154. doi:[10.24985/ijass.2017.29.2.143](https://doi.org/10.24985/ijass.2017.29.2.143)
37. Mulligan IJ, Biddington WB, Barnhart BD, Ellenbecker TS. Isokinetic profile of shoulder internal and external rotators of high school aged baseball pitchers. *J Strength Cond Res.* 2004;18(4):861-866. doi:[10.1519/14633.1](https://doi.org/10.1519/14633.1)
38. Pawlik D, Dziubek W, Rogowski Ł, Struzik A, Rokita A. Strength abilities and serve reception efficiency of youth female volleyball players. *Appl Bionics Biomech.* 2022;2022:4328761. doi:[10.1155/2022/4328761](https://doi.org/10.1155/2022/4328761)
39. Pontaga I, Zidens J. Shoulder rotator muscle dynamometry characteristics: side asymmetry and correlations with ball-throwing speed in adolescent handball players. *J Hum Kinet.* 2014;42:41-50. doi:[10.2478/hukin-2014-0059](https://doi.org/10.2478/hukin-2014-0059)
40. Saccol MF, Gracitelli GC, da Silva RT, et al. Shoulder functional ratio in elite junior tennis players. *Phys Ther Sport.* 2010;11(1):8-11. doi:[10.1016/j.ptsp.2009.11.002](https://doi.org/10.1016/j.ptsp.2009.11.002)

41. van Cingel R, Habets B, Willemsen L, Staal B. Shoulder dynamic control ratio and rotation range of motion in female junior elite handball players and controls. *Clin J Sport Med*. 2018;28(2):153-158. doi:[10.1097/jsm.0000000000000429](https://doi.org/10.1097/jsm.0000000000000429)
42. Vodička T, Zvonař M, Pačes J, Knjaz D, Ružbarský P, Zháněl J. Strength values of shoulder internal and external rotators in junior tennis players. *Kinesiology*. 2018;50(2):181-187. doi:[10.26582/k.50.2.9](https://doi.org/10.26582/k.50.2.9)
43. Yildiz Y, Aydin T, Sekir U, Kiralp MZ, Hazneci B, Kalyon TA. Shoulder terminal range eccentric antagonist/concentric agonist strength ratios in overhead athletes. *Scand J Med Sci Sports*. 2006;16(3):174-180. doi:[10.1111/j.1600-0838.2005.00471.x](https://doi.org/10.1111/j.1600-0838.2005.00471.x)
44. Bassett R, Browne A, Morrey B, An K. Glenohumeral muscle force and moment mechanics in a position of shoulder instability. *J Biomech*. 1990;23(5):405-415. doi:[10.1016/0021-9290\(90\)90295-E](https://doi.org/10.1016/0021-9290(90)90295-E)
45. Durall CJ, Davies GJ, Kernozek TW, Gibson MH, Fater DC, Straker JS. The effects of training the humeral rotators on arm elevation in the scapular plane. *J Sport Rehabil*. 2001;10(2):79-92. doi:[10.1123/jsr.10.2.79](https://doi.org/10.1123/jsr.10.2.79)
46. Ivey FM Jr, Calhoun JH, Rusche K, Bierschenk J. Isokinetic testing of shoulder strength: normal values. *Arch Phys Med Rehabil*. 1985;66(6):384-386.
47. Scoville CR, Arciero RA, Taylor DC, Stoneman PD. End range eccentric antagonist/concentric agonist strength ratios: a new perspective in shoulder strength assessment. *J Orthop Sports Phys Ther*. 1997;25(3):203-207. doi:[10.2519/jospt.1997.25.3.203](https://doi.org/10.2519/jospt.1997.25.3.203)
48. Tanaka N, Sakai Y, Maruyama Y, Hirayama T, Iwamoto W, Wagatsuma K. Throwing kinematics and elbow varus torque relative to ball size in junior baseball players. *J Phys Ther Sci*. 2022;34(1):22-25. doi:[10.1589/jpts.34.22](https://doi.org/10.1589/jpts.34.22)
49. Hirashima M, Yamane K, Nakamura Y, Ohtsuki T. Kinetic chain of overarm throwing in terms of joint rotations revealed by induced acceleration analysis. *J Biomech*. 2008;41(13):2874-2883. doi:[10.1016/j.jbiomech.2008.06.014](https://doi.org/10.1016/j.jbiomech.2008.06.014)
50. Pawlowski D, Perrin DH. Relationship between shoulder and elbow isokinetic peak torque, torque acceleration energy, average power, and total work and throwing velocity in intercollegiate pitchers. *Athl Train*. 1989;24(2):129-132.
51. Agresta CE, Krieg K, Freehill MT. Risk factors for baseball-related arm injuries: A systematic review. *Orthop J Sports Med*. 2019;7(2):2325967119825557. doi:[10.1177/2325967119825557](https://doi.org/10.1177/2325967119825557)
52. Nakaji RM, Ellenbecker TS, McClenahan KM, Roberts LM, Perez C, Dickenson SB. Descriptive strength and range of motion in youth baseball players. *Int J Sports Phys Ther*. 2021;16(1):195-206. doi:[10.26603/001c.18815](https://doi.org/10.26603/001c.18815)

SUPPLEMENTARY MATERIALS

Appendix A

Download: <https://ijspt.scholasticahq.com/article/125765-isokinetic-dynamometry-for-external-and-internal-rotation-shoulder-strength-in-youth-athletes-a-scoping-review/attachment/253027.docx>
