

Original Research

Are We Overlooking Anatomical Contributions to Dynamic Knee Valgus?

Matt Dewald^{1a}, Madison Andersen, Laura Higgins, Emma Porter, Alex Wickersham

¹ Physical Therapy, University of South Dakota

Keywords: dynamic knee valgus, jump performance, kinematics, bony anatomy, 2-D analysis.

<https://doi.org/10.26603/001c.128587>

International Journal of Sports Physical Therapy

Vol. 20, Issue 2, 2025

Background

Dynamic knee valgus (DKV) is widely considered a risk factor for injuries, despite contradictory research. Consequently, athletic performance and injury rehabilitation cueing has primarily focused on keeping the “knees out”.

Purpose

The purpose of this study was to assess jump performance measures and anatomical contributions.

Study Design

Cross-Sectional Study

Methods

Jump height, ground contact time, reactive strength index, and DKV was collected with the MyJump2 and Coach My Video apps. Static anatomical measurements were collected. Subjects completed nine jumps with each leg using the same set-up; performing three single leg six-inch depth jumps with their natural form, three depth jumps with external cuing towards increased DKV, and three depth jumps with cuing towards no DKV. ANOVA was used to compare jump data. Pearson Correlation Coefficients were used to assess relationships between DKV and anatomical measurements, jump height, ground contact time, and reactive strength index. Intraclass correlation coefficient (ICC) was used to assess inter-rater reliability of MyJump2 and Coach My Video measurements.

Results

50 subjects (35 included) participated in this study. With a cued DKV jump, ground contact time had a moderate positive correlation with DKV measurements ($r=.49$, $p<0.01$), however, this was not the case with subjects' natural jump and cued no DKV alignment jumps. Static anatomical measurements of static knee valgus and Q-angle had a weak positive correlation with DKV measurements for subjects' natural jumps ($r=.37$, $p<0.01$ and $r=.34$, $p=0.04$, respectively). When DKV measurements were normalized to an anatomical measurement, no correlations existed with any of the performance measurements. There was very strong inter-rater reliability ($ICC=.96-.99$) of all the measurements.

^a Corresponding author:
Matt Dewald, PT, DPT, OCS, SCS
University of South Dakota
Department of Physical Therapy
414 E. Clark St.
Vermillion, SD 57069
Phone: 605-658-6359
Fax: 605-638-5637
Email: matt.dewald@usd.edu

Conclusion

Bony anatomical alignment should be considered alongside kinematics, as normalization controlled for the differences in DKV. Future research should normalize DKV measurements by bony anatomy when addressing DKV and jump performance.

Level of Evidence

2c

INTRODUCTION

Dynamic knee valgus (DKV), also known as dynamic genu valgus is defined as a lower limb movement pattern that is a combination of internal rotation and adduction of the femur, knee abduction, anterior tibial translation, and external tibial rotation.¹ The normal range of DKV of a single-leg landing task has been reported as 1-9 degrees for males and 5-12 degrees for females when measured from a digital video camera two-dimensionally.² Anatomical characteristics contribute to DKV, with the Q-angle being the most significant.³⁻⁵ Q-angles are measured with similar, although different anatomical landmarks than knee valgus. DKV and static knee valgus are measured from the Anterior Superior Iliac Spine (ASIS) to the mid-patella to a line along the tibia between the medial and lateral malleoli. Meanwhile, Q-angle is measured from the ASIS, through the mid-patella, and ending on a line through the middle of the tibial tubercle.

Health professionals and coaches traditionally have considered DKV as being associated with increased risk of injuries and decreased jump performance. This has commonly led to a “knees out” cueing to prevent DKV during athletic performance, to the point that some clinicians attempt to avoid any degree of DKV. For example, the International Olympic Committee has published a review on ACL injuries encouraging clinicians to avoid excessive dynamic knee valgus and focus on a knee over toe position.⁶ Additionally, static anatomical measurements, such as Q-angle and Craig’s test, are rarely considered when providing feedback to athletes regarding their DKV.

Presently, there is contradictory evidence on DKV and injury risk.⁷⁻¹³ The 2005 prospective study by Hewett et al. identifying DKV as the primary culprit of ACL injury in female athletes has most commonly been referenced by clinicians as the reason to avoid DKV.¹⁰ This study reported those suffering an ACL injury (n=9) had 8 degrees more DKV than those who did not (n=196).¹⁰ Conversely with a similar set-up Nilstad et al. identified no association between DKV and ACL tears in female athletes who sustained an ACL injury (n=56) compared to uninjured athletes (n=722).⁷ When synthesizing the body of literature on DKV and ACL injuries, a 2020 meta-analysis of the nine highest quality articles with over 1300 total subjects found no association between DKV and ACL injuries.⁸

Additionally, many studies suggest a positive association between DKV and patellofemoral pain, stress fractures, iliotibial band syndrome, and Achilles’ tendinopathy. However, these studies have not considered the static bony anatomy of the subjects, overlooking the possible contributions of anatomical characteristics.³⁻⁵ In parallel, bony

anatomy of static knee valgus was found to have a significant relationship with DKV during a drop-jump, while foot alignment, hamstring strength, quadriceps strength, and hip abductor strength had no association.³ Furthermore, anatomical measurements in both the frontal and transverse planes at the hips and knees have been shown to have an association with DKV during double-leg landing, further adding value to the idea that anatomical differences contribute to dynamic measures.¹⁴


Moreover, there is conflicting evidence on the relationship between DKV and jump performance.^{5,15-17} Confounding variables such as specific training effects and bony anatomy was not considered with these reports. There is also dispute regarding the neuromuscular contributions to DKV, specifically, the association of hip abductor muscle weakness to DKV.^{5,18-22} Finally, interventional studies for improving DKV have had mixed results, with some studies demonstrating improved DKV with training proximally at the hip and core, while others found no change.^{23,24} Seeing that there is conflicting evidence surrounding the current consensus on DKV, additional information is needed to establish a better understanding of these relationships. The purpose of this study was to assess jump performance measures and anatomical contributions.

METHODS

STUDY DESIGN AND SUBJECTS

This cross-sectional study recruited 50 subjects from the university wellness center between the ages of 18 and 30 years old who felt they could complete nine single leg depth jumps from a 6-inch box with each leg. A depth jump consists of dropping from a box of a predetermined height and immediately, upon landing a single maximal jump is performed.²⁵ All subjects were recruited from a common area near the entrance of the wellness center. All data were collected at this location as well. The subjects were incentivized with a sports drink for their participation. Fifteen subjects were excluded following poor execution of a cued jump for dynamic knee valgus, primarily due to turning their foot inward. After consenting to the research, background demographic information including sex, age, weight, height, injury history, and past sport participation was collected from subjects. BMI was later calculated based on the subjects’ self-reported height and weight.

Static anatomical measurements of transverse plane femoral anteversion and frontal plane genu valgus were completed by the same researcher. Femoral anteversion was measured with the Craig’s test, while genu valgus was measured with Q-angle²⁶ and static knee valgus. Craig’s test is



A. B. C.
Figure 1. Subject performing the single-leg depth jump. A. The subject is demonstrating their natural form. B. The subject is demonstrating a DKV cued depth jump. C. The subject is demonstrating a no DKV cued depth jump.

the most commonly used clinical assessment of femoral anteversion, however, the validity of this test has been questioned.²⁷ Souza and Powers reported good reliability with a strong positive correlation ($r=.61$, $p<0.01$) when compared with MRI measurements of femoral anteversion.²⁸ This suggests that Craig's test can be a useful clinical test for screening femoral anteversion without higher costs associated with imaging.

Static knee valgus and 2-Dimensional DKV utilize the same landmarks: from the subject's ASIS, through the mid-patella, and ending at the midline of the anterior tibia between the medial and lateral malleoli. Q-angle was measured from the ASIS, through the mid-patella, and ending on a line through the middle of the tibial tubercle. Both the Q-angle and static knee valgus were measured in supine. Landmarks were marked with either tape or marker to facilitate recording of DKV. The sample size was determined using G*Power Version 3.1.9.7 with .05 alpha, .8 power, and a medium-large effect size of .5 resulting in 34 subjects needed, thus, 50 were recruited in anticipation of dropouts and excluded subjects.

VARIABLES

The independent variables were the three single-leg depth jump modes: a natural jump form, jump form with cued DKV, and jump form cued for no DKV. The natural depth jump (Figure 1a) consisted of the subjects jumping with no cue. The DKV cued depth jump (Figure 1b) consisted of a "knees-in" verbal cue with an external reference cone 30 degrees medial. The no DKV cued depth jump (Figure 1c) consisted of a verbal "knees straight forward" cue with an external reference cone directly in front of the landing zone. A single-leg depth jump was chosen over a double-leg, due to the greater amount of knee valgus that often occurs with single-leg.²⁹ The different jump forms were chosen to assess changes in jump performance based on DKV angles.

The dependent variables measured were jump height, ground contact time, reactive strength index (RSI), and DKV of the jump. Additionally, a true DKV measurement, normalized for frontal plane anatomical measurements, was collected consisting of subtracting the subject's static knee valgus from their DKV measurements. True DKV was calculated as it represents the degree of frontal plane motion that was occurring, controlling for anatomical differences between subjects. True DKV provides insights into DKV motor control. Three trials were collected for each leg and each jump with means being recorded for each subject. Static anatomical measurements included Craig's test for femoral anteversion, Q-angle for genu valgus, and static knee valgus for genu valgus.²⁶ Static knee valgus was used in calculating true DKV measurement as it was the most ecologically similar to DKV and it had the highest and strongest correlation with DKV ($r=.37$, $p=0.03$).

DATA SOURCES/MEASUREMENTS

The MyJump2 application was used to measure ground reaction time, RSI, and jump height. MyJump2 is a reliable and valid tool, frequently used in jump studies.³⁰⁻³² The Coach My Video application was used to measure DKV. DKV was measured at the most extreme point, often at the deepest point of the depth jump. 2-D phone-based apps have been found reliable and validated for assessing kinematics.³³⁻³⁵ The apps were installed on a university issued iPad.

A 6-inch box was used for the depth jump with an iPad placed seven feet in front of the box with the camera mounted 15 inches from the ground to standardize recordings of the jumps. To ensure focus on jump performance, subjects were told to jump as high and as quickly as possible. Each jump was demonstrated by the same researcher for every subject. The subjects were cued with "Point your knee towards the cone." Subjects were given the option to perform a slow, step-through practice trial as well as full-speed practice trials. Subjects were given as many practice

trials as they felt necessary to gain full understanding of the test. All subjects began with their natural jump form ([Figure 1a](#)), completing three jumps with their right and then three on their left. Random numbers were used to determine whether the cued DKV ([Figure 1b](#)) or cued no DKV ([Figure 1c](#)) would be the second or third set of jumps. The same procedure of completing three jumps for the right leg was completed before switching to the left for the cued jumps.

BIAS

To ensure reliability of anatomical characteristics, the same researcher placed markers on designated landmarks and measured Q-angle, static knee valgus, and Craig's test for all sessions. Craig's test validity is not that of CT or MRI, although it is regarded as an acceptable clinical exam to screen for femoral anteversion in the absence of imaging.^{27, 28} Therefore, transverse plane anatomical contributions to DKV should be interpreted cautiously. Randomization of the DKV and no DKV jumps was used to reduce order effects. Performance bias was controlled with a standardized assessment preserving fidelity of the protocol. The high reliability of the measurement techniques used to assess jump performance and DKV ensures quality of the data. Two blinded researchers not associated with the study assessed the jump performances and kinematics at a different and neutral site, minimizing expectation bias. There was no missing data.

STATISTICAL METHODS

All statistical analysis was completed with SAS. All alphas were set at 0.05. Normality of the jump variables were assessed with Shapiro-Wilk, Histograms, and Q-Q plots.³⁶ ANOVA was used to compare jump height, contact time, RSI, and DKV with each jump method. Pearson Correlation Coefficients were used to assess DKV and anatomical measurements, jump height, ground contact time, and RSI. Intraclass correlation coefficient (ICC) was used to assess inter-rater reliability of the measurement tools. Strength of correlations was described using the following scale outlined by the British Medical Journal: 0-0.19 is regarded as very weak, 0.2-0.39 as weak, 0.40-0.59 as moderate, 0.6-0.79 as strong, and 0.8-1 as very strong correlation.³⁷

RESULTS

SUBJECTS

Fifty subjects were recruited and completed the study. Following analysis of the jump data, 15 subjects' data were excluded as they did not complete the cued DKV jump correctly, pointing their feet too far towards midline. The remaining 35 subjects were included in final data analysis.

DESCRIPTIVE DATA

The average age of subjects was 21.91 years-old (95% CI 21.12-22.71). Their average height was 68.97 inches (95%

Table 1. Subject characteristics, reported as mean (95% CI), unless otherwise indicated

Variable	Outcome
Number of Subjects, n	35
Females, n (%)	18 (51.4%)
Age (years)	21.91 (21.12-22.71)
Weight (pounds)	165.94 (154.93-176.96)
Height (inches)	68.97 (67.81-70.13)
BMI (kg/m ²)	24.38 (23.15-25.61)
Q-angle (degrees)	16.26 (14.86-17.66)
Static knee valgus (degrees)	9.90 (8.94-10.86)
Craig's Test (degrees)	9.40 (8.57-10.23)
Significant Injury History, n (%)	15 (42.9%)
Past Physical Therapy, n (%)	18 (51.4%)
Jump Training History, n (%)	27 (77.1%)

Abbreviations: n= number of subjects, CI= Confidence Interval, BMI= Body Mass Index, kg= Kilogram, m= Meter, Q-angle= Quadriceps Angle

CI 67.81-70.13). Their average weight was 165.94 pounds (95% CI 154.93-176.96). Their average BMI was 24.38 (95% CI 23.15-25.61). Their average Q-angle was 16.26 degrees (95% CI 14.86-17.66). Static knee valgus was 9.90 degrees (95% CI 8.94-10.86). Craig's test was 9.40 degrees (95% CI 8.57-10.23). Over half (51.4%) of subjects had past physical therapy, 42.9% had a history of significant lower extremity injury, and 77.1% had past jump training. All subject characteristics can be found in [Table 1](#).

When stratifying the study characteristics by sex, five variables were significantly different between females and males. Weight, height, BMI, Q-angle, and static knee valgus were all statistically different. See [Table 2](#) for subject characteristics stratified by sex.

OUTCOME DATA

The average normal single leg depth jump height was 5.87 inches (95% CI 4.99-6.76). The average contact time was 419.72 ms (95% CI 388.71-450.73). The average RSI was .86 (95% CI .76-.97). The average DKV with the subjects' natural jump was 13.12 degrees (95% CI 11.29-14.96). The true DKV was 3.22 degrees (95% CI 1.50-4.95). [Table 3](#) contains the jump performance data of the included subjects.

Jump height and DKV were significantly different when comparing the jump performances based on sex. Interestingly, there were no statistically significant differences for DKV between sexes once the measure was normalized for static frontal plane anatomical measures, also known as the *true DKV*. See all subject jump characteristics stratified by sex in [Table 4](#).

MAIN RESULTS

Performance and DKV averages for each jump type can be found in [Table 5](#). There was no difference in jump heights between jump types ($p=0.59$). There was a significant difference between contact times, with the cuing for DKV jumps being longer than the natural ($p=0.03$) and cuing for no DKV jumps ($p=0.04$). RSI was significantly lower for the jump cued for DKV compared to the natural jump ($p=0.03$). Cuing for DKV was effective, as those jumps had a signifi-

Table 2. Subject characteristics stratified by sex, reported as mean (95% CI)

Variable	Subjects	Female	Male	p value*
Number of Subjects, n	35	18	17	-
Age (years)	21.91 (21.12-22.71)	22.44 (21.27-23.62)	21.35 (20.23-22.47)	0.16
Weight (pounds)	165.94 (154.93-176.96)	151.11 (135.56-166.67)	181.65 (168.69-194.60)	<0.01
Height (inches)	68.97 (67.81-70.13)	67.17 (65.46-68.87)	70.88 (69.82-71.94)	<0.01
BMI (kg/m ²)	24.38 (23.15-25.61)	23.40 (21.63-25.16)	25.41 (23.66-27.16)	0.01
Q-angle (degrees)	16.26 (14.86-17.66)	17.75 (15.61-19.89)	14.68 (13.01-16.34)	0.02
Static knee valgus (degrees)	9.90 (8.94-10.86)	11.58 (10.36-12.81)	8.12 (7.17-9.06)	<0.01
Craig's Test (degrees)	9.40 (8.57-10.23)	9.67 (8.63-10.70)	9.12 (7.69-10.54)	0.51

Abbreviations: n= number of subjects, CI= Confidence Interval, BMI= Body Mass Index, kg= Kilogram, m= Meter, Q-angle= Quadriceps Angle

*p values reference differences between female and male subjects

Table 3. Subject jump characteristics, reported as mean (95% CI)

Variable	Outcome
Jump Height (inches)	5.87 (4.99-6.76)
Contact Time (ms)	419.72 (388.71-450.73)
RSI	.86 (.76-.97)
DKV (degrees)	13.12 (11.29-14.95)
True DKV (degrees)	3.22 (1.50-4.95)

Abbreviations: CI= Confidence Interval, ms= milliseconds, RSI= Reactive Strength Index, DKV= Dynamic Knee Valgus

Table 4. Subject jump characteristics stratified by sex, reported as mean (95% CI)

Variable	Included Subjects	Female	Male	p value*
Jump Height (inches)	5.87 (4.99-6.76)	4.53 (3.74-5.32)	7.29 (5.90-8.69)	<0.01
Contact Time (ms)	419.72 (388.71-450.73)	422.37 (376.11-468.63)	416.92 (370.63-463.20)	0.86
RSI	.86 (.76-.97)	.78 (.63-.92)	.95 (.81-1.10)	0.08
DKV (degrees)	13.12 (11.29-14.95)	14.97 (12.53-17.41)	11.16 (8.49-13.83)	0.03
True DKV (degrees)	3.22 (1.50-4.95)	3.39 (1.07-5.71)	3.04 (.21-5.88)	0.84

Abbreviations: CI= Confidence Interval, ms= milliseconds, RSI= Reactive Strength Index, DKV= Dynamic Knee Valgus, *p values reference female and male subjects

Table 5. Performance and Kinematic Measurements of Jump Types, reported as mean (95% CI)

	Natural	DKV	No DKV
Jump Height (inches)	5.87 (4.99-6.76)	5.38 (4.52-6.23)	5.98 (5.04-6.92)
Contact Time (ms)	419.72 (388.71-450.73)	528.47 (467.93-589.02)	448.63 (408.10-489.15)
RSI	.86 (.76-.97)	0.68 (.59-.77)	.82 (.72-.92)
DKV (Degrees)	13.12 (11.29-14.95)	35.02 (31.52-38.52)	13.79 (12.03-15.55)
True DKV (Degrees)	3.22 (1.50-4.95)	25.12 (21.71-28.53)	3.89 (2.13-5.64)

Abbreviations: CI= Confidence Interval, ms= milliseconds, RSI= Reactive Strength Index, DKV= Dynamic Knee Valgus

cantly higher DKV ($p < 0.01$) than the natural jump and cutting for no DKV. Interestingly, the jumps in which subjects were cued for no DKV did not lower the DKV below the subjects' natural jumps.

Craig's test was not correlated to DKV measurements of natural jumps ($p = 0.72$). Both static knee valgus ($r = .37$, $p = 0.03$) and Q-angle ($r = .34$, $p = 0.04$) had positive, weak significant correlations with DKV measurements for the subjects' natural jumps. Static knee valgus demonstrated a greater positive significant correlation with DKV than the traditional Q-angle and the adjusted R-square suggests that the static knee valgus accounts for 11% of the DKV (Figure 2).

The ICC for jump height, ground contact time, and RSI measurements taken with the MyJump2 app was near per-

fect at .99 for each. The ICC for DKV measured with the Coach My Video app was also very strong at .96.

DISCUSSION

This study highlights the need to consider static bony alignment when assessing DKV with jump performances, as the differences in DKV between sexes was accounted for by the frontal plane anatomy. Poor clinical decision making can occur when making assumptions about movement faults without measuring anatomical contributions. Furthermore, static knee valgus could be considered over the use of Q-angles since using the same landmarks for static and dynamic tasks increases the ecological validity of applying static measurements. That being so, the static knee




Figure 2. Correlation of DKV with static knee valgus.

Table 6. Inter rater reliability of dependent variables

	ICC
Jump Height	.99
Ground Contact Time	.99
Reactive Strength Index	.99
Dynamic Knee Valgus	.96

Abbreviations: ICC= Intraclass Correlation Coefficients

valgus measurements were more highly correlated to DKV in this study.

Since the no DKV cued jump did not lower the DKV of subjects, in fact, increased it, the researchers question the effectiveness of using an external visual cue. When cued for DKV, ground contact time had a significant correlation with true DKV measurements, suggesting that when a patient has high DKV, instead of cuing to keep the knee from caving in, clinicians could cue for less contact time to indirectly control DKV. While external cuing has demonstrated effectiveness, it traditionally has not focused on contact time for improving DKV.³⁸⁻⁴⁰ Motor learning studies have found that cuing for contact time has improved both ground contact time and RSI, however, these studies have not assessed the impact on DKV.^{41,42} RSI was significantly lower for the jump cued for DKV compared to the natural jump ($p=0.03$), likely related to the contribution of contact time on RSI. However, increased impulse and impact forces have been associated with cueing for less ground contact time, both of which have been linked to increased injury risk.^{41,43,44}

When considering improving jump height, this study did not find any difference within subjects when considering DKV. This is comparable to findings on squat jumps and

countermovement jumps in past research.¹⁶ This study's findings were contrary to a 3-month interventional study that found improved jump height with an exercise program targeting less DKV.¹⁷ However, that interventional study only had eight subjects complete the intervention and did not control for confounding that the improvements in jump height could be linked to improved strength and power from the exercise program and not decreased DKV.¹⁷

The high reliability of the apps found in this study is encouraging for clinicians without an expensive performance laboratory. These apps are readily accessible and do not require extensive time and monetary investments. When considering this along with the high reliability and validity of 2-D phone apps found in past studies, clinicians can be confident in the reliability of using app-based 2-D analysis tools for assessing jump performance and DKV.³⁰⁻³⁵

It must be noted that the population used in this study was primarily college students, but that does not change the fact that their anatomical alignment contributes to their DKV. These considerations are most dramatic for females, as females typically have increased frontal plane knee valgus and femoral anteversion, both of which can masquerade as DKV if not assessed.⁴ Additionally, the validity of craig's test has been questioned, potentially over or underrepresenting transverse plane anatomical contributions in this study. Thirty percent of subjects were excluded due to not being able to complete the cued DKV jump correctly, these subjects could have swayed the results or could have added statistical power if included. Additionally, the fact that 30 percent of the subjects struggled with the cued DKV jump suggests difficulty in completing the task and possibly less validity when comparing to natural jumping mechanics. Participants who had experienced past injuries, physical therapy, and jump training were not excluded from

this study. The influence of previous rehab and jump training may have impacted performance throughout this study. Additionally, the external cuing may have caused subjects to focus more on the cone rather than their best jump performance.

CONCLUSION

Bony alignment should be considered with kinematics, as normalization of DKV measurements by static anatomical measurements controlled for the differences in DKV between men and women during subjects' natural jumping performance. When jumping with cued DKV alignment, ground contact time had a significant positive correlation

with DKV measurements, therefore cueing for less ground contact time may be an external cue to use to control for excessive DKV, although prospective studies would be needed to test this in the future.

.....

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

Submitted: July 05, 2024 CST. Accepted: December 14, 2024 CST. Published: February 01, 2025 CST.

© The Author(s)



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

REFERENCES

1. Wilczyński B, Zorena K, Ślęzak D. Dynamic knee valgus in single-leg movement tasks. Potentially modifiable factors and exercise training options. A literature review. *Int J Environ Res Public Health*. 2020;17(21):8208. doi:[10.3390/ijerph17218208](https://doi.org/10.3390/ijerph17218208)
2. Herrington L, Munro A. Drop jump landing knee valgus angle; normative data in a physically active population. *Phys Ther Sport*. 2010;11(2):56-59. doi:[10.1016/j.ptspt.2009.11.004](https://doi.org/10.1016/j.ptspt.2009.11.004)
3. Nilstad A, Krosshaug T, Kam-Ming MOK, Bahr R, Andersen TE. Association between anatomical characteristics, knee laxity, muscle strength, and peak knee valgus during vertical drop-jump landings. *J Ortho Sports Ther*. 2015;45(12):998-1005. doi:[10.2519/jospt.2015.5612](https://doi.org/10.2519/jospt.2015.5612)
4. Almeida GP, Silva AP, França FJ, Magalhães MO, Burke TN, Marques AP. Relationship between frontal plane projection angle of the knee and hip and trunk strength in women with and without patellofemoral pain. *J Back Musculoskelet Rehabil*. 2016;29(2):259-266. doi:[10.3233/BMR-150622](https://doi.org/10.3233/BMR-150622)
5. Stickler L, Finley M, Gulgin H. Relationship between hip and core strength and frontal plane alignment during a single leg squat. *Phys Ther Sport*. 2015;16(1):66-71. doi:[10.1016/j.ptspt.2014.05.002](https://doi.org/10.1016/j.ptspt.2014.05.002)
6. Renstrom P, Ljungqvist A, Arendt E, et al. Non-contact ACL injuries in female athletes: an International Olympic Committee current concepts statement. *Br J Sports Med*. 2008;42(6):394-412. doi:[10.1136/bjsm.2008.048934](https://doi.org/10.1136/bjsm.2008.048934)
7. Nilstad A, Petushek E, Mok KM, Bahr R, Krosshaug T. Kiss goodbye to the “kissing knees”: no association between frontal plane inward knee motion and risk of future non-contact ACL injury in elite female athletes. *Sports Biomech*. 2023;22(1):65-79. doi:[10.1080/14763141.2021.1903541](https://doi.org/10.1080/14763141.2021.1903541)
8. Cronström A, Creaby MW, Ageberg E. Do knee abduction kinematics and kinetics predict future anterior cruciate ligament injury risk? A systematic review and meta-analysis of prospective studies. *BMC Musculoskelet Disord*. 2020;21(1):563. doi:[10.1186/s12891-020-03552-3](https://doi.org/10.1186/s12891-020-03552-3)
9. Goerger BM, Marshall SW, Beutler AI, Blackburn JT, Wilckens JH, Padua DA. Anterior cruciate ligament injury alters preinjury lower extremity biomechanics in the injured and uninjured leg: the JUMP-ACL study. *Br J Sports Med*. 2015;49(3):188-195. doi:[10.1136/bjsports-2013-092982](https://doi.org/10.1136/bjsports-2013-092982)
10. Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med*. 2005;33(4):492-501. doi:[10.1177/0363546504269591](https://doi.org/10.1177/0363546504269591)
11. Krosshaug T, Steffen K, Kristianslund E, et al. The vertical drop jump is a poor screening test for ACL injuries in female elite soccer and handball players: a prospective cohort study of 710 athletes. *Am J Sports Med*. 2016;44(4):874-883. doi:[10.1177/0363546515625048](https://doi.org/10.1177/0363546515625048)
12. Leppanen M, Pasanen K, Kujala UM, et al. Stiff landings are associated with increased ACL injury risk in young female basketball and floorball players. *Am J Sports Med*. 2017;45(2):386-393. doi:[10.1177/0363546516665810](https://doi.org/10.1177/0363546516665810)
13. Numata H, Nakase J, Kitaoka K, et al. Two-dimensional motion analysis of dynamic knee valgus identifies female high school athletes at risk of non-contact anterior cruciate ligament injury. *Knee Surg Sports Traumatol Arthrosc*. 2018;26(2):442-447. doi:[10.1007/s00167-017-4681-9](https://doi.org/10.1007/s00167-017-4681-9)
14. Uota S, Nguyen AD, Aminaka N, Shimokochi Y. Relationship of knee motions with static leg alignments and hip motions in frontal and transverse planes during double-leg landing in healthy athletes. *J Sport Rehabil*. 2017;26(5):396-405. doi:[10.1123/jsr.2016-0053](https://doi.org/10.1123/jsr.2016-0053)
15. Arundale AJH, Kvist J, Häggglund M, et al. Jump performance in male and female football players. *Knee Surg Sports Traumatol Arthrosc*. 2020;28:606-613. doi:[10.1007/s00167-019-05747-1](https://doi.org/10.1007/s00167-019-05747-1)
16. Lagares LS, de Silva MS, de Macedo R, et al. Dynamic knee valgus and its relationship with performance in the countermovement jump and the squat jump. *J Ex Phys*. 2021;24(6):1-10.
17. Giustino V, Messina G, Patti A, et al. Effects of a postural exercise program on vertical jump height in young female volleyball players with knee valgus. *Int J Environ Res Public Health*. 2022;19(7):3953. doi:[10.3390/ijerph19073953](https://doi.org/10.3390/ijerph19073953)
18. Hollman JH, Ginos BE, Kozuchowski J, Vaughn AS, Krause DA, Youdas JW. Relationships between knee valgus, hip-muscle strength, and hip-muscle recruitment during a single-limb step-down. *J Sport Rehabil*. 2009;18(1):104-117. doi:[10.1123/jsr.18.1.104](https://doi.org/10.1123/jsr.18.1.104)

19. Baldon R de M, Piva SR, Scattone Silva R, Serrão FV. Evaluating eccentric hip torque and trunk endurance as mediators of changes in lower limb and trunk kinematics in response to functional stabilization training in women with patellofemoral pain. *Am J Sports Med.* 2015;43(6):1485-1493. doi:[10.1177/0363546515574690](https://doi.org/10.1177/0363546515574690)
20. Bredeweg S, Buist I. No relationship between running related injuries and kinetic variables. *Br J Sports Med.* 2011;45:328. doi:[10.1136/bjsm.2011.084038.52](https://doi.org/10.1136/bjsm.2011.084038.52)
21. Dicharry J. Kinematics and kinetics of gait: from lab to clinic. *Clin Sports Med.* 2010;29(3):347-364. doi:[10.1016/j.csm.2010.03.013](https://doi.org/10.1016/j.csm.2010.03.013)
22. Llurda-Almuzara L, Pérez-Bellmunt A, López-de-Celis C, et al. Normative data and correlation between dynamic knee valgus and neuromuscular response among healthy active males: a cross-sectional study. *Sci Rep.* 2020;10:17206. doi:[10.1038/s41598-020-74177-8](https://doi.org/10.1038/s41598-020-74177-8)
23. Baldon R, Serrao FV, Silva RS, Piva SR. Effects of functional stabilization training on pain, function, and lower extremity biomechanics in women with patellofemoral pain: A randomized clinical trial. *J Orthop Sports Phys Ther.* 2014;44(4):240-251. doi:[10.2519/jospt.2014.4940](https://doi.org/10.2519/jospt.2014.4940)
24. Mozafaripour E, Seidi F, Minoonejad H, Bayattork M, Khoshroo F. The effectiveness of the comprehensive corrective exercise program on kinematics and strength of lower extremities in males with dynamic knee valgus: a parallel-group randomized wait-list controlled trial. *BMC Musculoskelet Disord.* 2022;23(1):700. doi:[10.1186/s12891-022-05652-8](https://doi.org/10.1186/s12891-022-05652-8)
25. Radcliffe J. Form and safety in plyometric training. *Perf Train J.* 2003;2(2):21-25.
26. Horton MG, Hall TL. Quadriceps femoris muscle angle: normal values and relationships with gender and selected skeletal measures. *Phys Ther.* 1989;69(11):897-901. doi:[10.1093/ptj/69.11.897](https://doi.org/10.1093/ptj/69.11.897)
27. Uding A, Bloom NJ, Commean PK, et al. Clinical tests to determine femoral version category in people with chronic hip joint pain and asymptomatic controls. *Musculoskelet Sci Pract.* 2019;39:115-122. doi:[10.1016/j.msksp.2018.12.003](https://doi.org/10.1016/j.msksp.2018.12.003)
28. Souza RB, Powers CM. Concurrent criterion-related validity and reliability of a clinical test to measure femoral anteversion. *J Orthop Sports Phys Ther.* 2009;39(8):586-592. doi:[10.2519/jospt.2009.2996](https://doi.org/10.2519/jospt.2009.2996)
29. Neal BS, Lack SD, Lankhorst NE, Raye A, Morrissey D, van Middelkoop M. Risk factors for patellofemoral pain: A systematic review and meta-analysis. *Br J Sports Med.* 2019;53(5):270-281. doi:[10.1136/bjsports-2017-098890](https://doi.org/10.1136/bjsports-2017-098890)
30. Alias MF, Hasan H, Miswan MS, Man DD. Inter-rater reliability and intra-rater reliability testing of my jump 2 mobile application in measuring counter movement jump. *Int J Health Move Sci.* 2021;9(6):1319-1323. doi:[10.13189/saj.2021.090628](https://doi.org/10.13189/saj.2021.090628)
31. Bogataj Š, Pajek M, Hadžić V, Andrašić S, Padulo J, Trajković N. Validity, reliability, and usefulness of My Jump 2 app for measuring vertical jump in primary school children. *Int J Environ Res Public Health.* 2020;17(10):3708. doi:[10.3390/ijerph17103708](https://doi.org/10.3390/ijerph17103708)
32. Haynes T, Bishop C, Antrobus M, Brazier J. The validity and reliability of the My Jump 2 app for measuring the reactive strength index and drop jump performance. *J Sports Med Phys Fit.* 2019;59(2):253-258. doi:[10.23736/S0022-4707.18.08195-1](https://doi.org/10.23736/S0022-4707.18.08195-1)
33. McLean SG, Walker K, Ford KR, Myer GD, Hewett TE, van den Bogert AJ. Evaluation of a two dimensional analysis method as a screening and evaluation tool for anterior cruciate ligament injury. *Br J Sports Med.* 2005;39(6):355-362. doi:[10.1136/bjsm.2005.018598](https://doi.org/10.1136/bjsm.2005.018598)
34. Mousavi SH, Hijmans JM, Moeini F, et al. Validity and reliability of a smartphone motion analysis app for lower limb kinematics during treadmill running. *Phys Ther Sport.* 2020;43:27-35. doi:[10.1016/j.ptsp.2020.02.003](https://doi.org/10.1016/j.ptsp.2020.02.003)
35. Ortiz A, Rosario-Canales M, Rodriguez A, Seda A, Figueroa C, Venegas-Rios H. Reliability and concurrent validity between two-dimensional and three-dimensional evaluations of knee valgus during drop jumps. *J Sports Med.* 2016;7:65-73. doi:[10.2147/OAISM.S100242](https://doi.org/10.2147/OAISM.S100242)
36. Vittinghoff E, Glidden DV, Shiboski SC, McCulloch AE. *Regression Methods in Biostatistics: Linear, Logistic, Survival, and Repeated Measures Models.* 2nd ed. Springer; 2012. doi:[10.1007/978-1-4614-1353-0](https://doi.org/10.1007/978-1-4614-1353-0)
37. BMJ. Statistics at square one: 11. Correlation and regression. BMJ. Accessed January 19, 2025. <https://www.bmj.com/about-bmj/resources-readers/publications/statistics-square-one/11-correlation-and-regression>

38. Gokeler A, Benjaminse A, Hewett TE, et al. Feedback techniques to target functional deficits following anterior cruciate ligament reconstruction: implications for motor control and reduction of second injury risk. *Sports Med.* 2013;43(11):1065-1074. doi:[10.1007/s40279-013-0095-0](https://doi.org/10.1007/s40279-013-0095-0)
39. Gokeler A, Benjaminse A, Welling W, Alferink M, Eppinga P, Otten B. The effects of attentional focus on jump performance and knee joint kinematics in patients after ACL reconstruction. *Phys Ther Sport.* 2015;16(2):114-120. doi:[10.1016/j.ptsp.2014.06.002](https://doi.org/10.1016/j.ptsp.2014.06.002)
40. Myer GD, Ford KR, Brent JL, Hewett TE. An integrated approach to change the outcome part II: targeted neuromuscular training techniques to reduce identified ACL injury risk factors. *J Strength Cond Res.* 2012;26(8):2272-2292. doi:[10.1519/JSC.0b013e31825c2c7d](https://doi.org/10.1519/JSC.0b013e31825c2c7d)
41. Oliver JL, Barillas SR, Lloyd RS, Moore I, Pedley J. External cueing influences drop jump performance in trained young soccer players. *J Strength Cond Res.* 2021;35(6):1700-1706. doi:[10.1519/JSC.0000000000002935](https://doi.org/10.1519/JSC.0000000000002935)
42. Barker L, Bankers S, Farmer B, Siedlik J, Harry JR, Grindstaff T. The influence of verbal cues on drop jump landing strategies in NCAA Division I soccer players. *Am J Sports Sci.* 2021;9(2):37-42. doi:[10.11648/j.ajss.20210902.12](https://doi.org/10.11648/j.ajss.20210902.12)
43. Bates NA, Ford KR, Myer GD, Hewett TE. Timing differences in the generation of ground reaction forces between the initial and secondary landing phases of the drop vertical jump. *Clin Biomech (Bristol, Avon).* 2013;28(7):796-799. doi:[10.1016/j.clinbiomech.2013.07.004](https://doi.org/10.1016/j.clinbiomech.2013.07.004)
44. Krosshaug T, Nakamae A, Boden BP, et al. Mechanisms of anterior cruciate ligament injury in basketball: video analysis of 39 cases. *Am J Sports Med.* 2007;35(3):359-367. doi:[10.1177/0363546506293899](https://doi.org/10.1177/0363546506293899)